

Circulating Microparticle Double-Stranded Deoxyribonucleic Acid in Systemic Lupus Erythematosus

Dear Editor,

Systemic lupus erythematosus (SLE) is a systemic autoimmune disease with complex pathogenesis which is yet to be completely understood. Since the original discovery of anti-double-stranded deoxyribonucleic acid (dsDNA) antibodies, the general result from intense research has been that native mammalian dsDNA is not immunogenic, even if presented with complete Freund's adjuvant.¹⁻³ This is despite an antigen-driven response as evidenced by the clonality of anti-dsDNA antibodies and patterns of random somatic mutations in both patients and murine models of SLE to suggest that DNA is the selecting antigen.^{2,4} These studies demonstrate that mechanisms for the production of anti-dsDNA antibodies and their affinity maturation toward anti-dsDNA specificity are operational in SLE patients.⁴ The question that arises is what transforms DNA to the immunogenic form in patients with SLE. Microparticles (MPs) represent a heterogeneous population of membrane-bound vesicles with a diameter of 0.1 μm to 1 μm that are released by the budding of plasma membrane and express antigens specific of their parental origin.⁵ The release of MPs by various cell types is a ubiquitous process that gets accelerated during cellular activation and apoptosis.⁵ Upon initiation, translocation of phosphatidylserine from the inner to outer surface leaflet of the plasma membrane results in loss of normal phospholipid asymmetry.⁵ As shown in cell lines undergoing in vitro apoptosis, DNA is sequestered into granules and then packaged into separate apoptotic bodies.⁶ Whereas the mechanism mediating this cellular rearrangement is not known, the end result is the repositioning of nuclear constituents in a form that may be more accessible to the immune system.⁷ We hypothesise that in this model, MPs may provide a framework to intensify the immunogenicity of the component DNA to induce anti-dsDNA antibody production and therefore, the aim of the present study was to evaluate the putative role of DNA within MPs (MP DNA) as the circulating antigenic target.

The study was cross-sectional in design. Fourteen unselected patients (12 women and 2 men) fulfilling at least 4 of the American College of Rheumatology 1997 revised classification criteria for SLE were included.⁸ Disease activity was assessed using the Safety of Estrogens in Lupus Erythematosus National Assessment (SELENA)-modified SLE Disease Activity Index (SLEDAI).⁹ Eight healthy female individuals were included as control subjects.

Anti-dsDNA immunoglobulin G (IgG) titers, measured using indirect enzyme-linked immunosorbent assay (Bio-Rad, Hercules, CA, US), and other disease markers were performed by the clinical laboratory of National University Hospital. Circulating MPs were obtained from platelet-free plasma (PFP) obtained by successive centrifugations of venous blood. PicoGreen (Invitrogen, Waltham, MA, US), a membrane-permeable dsDNA specific dye, was used to label the MP samples. We analysed the samples on FACSCalibur flow cytometer (BD Biosciences, Franklin Lakes, NJ, US). An initial MP-size gate was set with the help of calibrating fluorescent 0.22 μm , 0.45 μm , 0.88 μm and 1.34 μm polystyrene beads (Spherotech, Lake Forest, IL, US). MP DNA were enumerated on the SSC/FL1 plot and defined as events that were sensitive to differential detergent lysis using 0.05% Triton X-100. Therefore, each detergent-sensitive FL1-positive event on the SSC/FL1 plot was considered as 1 MP DNA. Plasma concentrations (MP DNA/ μL) were calculated according to the actual flow rate of the flow cytometer, MP DNA count per unit of time and net dilution during sample preparation of the analysed sample. No distinct population of detergent-sensitive MPs on the SSC/FSC plot could be gated to measure total MP numbers. Some samples were treated with 1 U of RNase-free DNase (Promega, Fitchburg, WI, US) for 20 min at 37°C to assess sensitivity of MP DNA to DNase I.

We identified MP DNA in PFP, based on fluorescence and detergent sensitivity (Figs. 1a and 1b). With this approach, extracellular DNA (plasma DNA) not associated with MPs were also detected in PFP. Importantly, although plasma DNA remained intact after detergent treatment, MPs containing DNA were detergent-soluble, establishing their phospholipid composition. To evaluate if MP DNA is resistant to DNase activity, MPs were treated with DNase I, counterstained with PicoGreen and then analysed using flow cytometry. There was a reduction of MP DNA count following DNase I treatment (Fig. 1c). Further treatment with 0.05% Triton X-100 led to complete solubilisation of MP DNA by detergent (Fig. 1d). Thus, DNA packaged within MPs is protected from DNase activity. The median concentration of MP DNA/ μL in PFP of the 14 SLE patients was significantly higher than the 8 healthy controls (2460.28 [Q1; Q3 1010.91; 3416.26] vs 403.73 [Q1; Q3 222.26; 1801.88]; $P = 0.020$) (Fig. 2). The median age of the SLE patients was 39.5 (Q1; Q3 25.0; 54.8), with a median disease




Fig. 1. Demonstration of MP DNA through representative flow cytometry analysis in plasma from an SLE patient. Using PicoGreen labelling, 2 distinct populations of FITC+ nanoparticles were obtained. The more granular population was sensitive to 0.05% Triton X-100 detergent, indicating that this population of FITC+ nanoparticles were MPs with membrane phospholipids (MP DNA count = 9203 in Fig. 1a and MP DNA count = 806 in Fig. 1b). Using DNase I, the MP DNA population decreased (MP DNA count = 1686 in Fig. 1c), demonstrating that DNA packaged within MPs is protected from DNase activity. With the addition of 0.05% Triton X-100, the MP DNA population which was resistant to DNase I is now solubilised (MP DNA count = 420 in Fig. 1d).

duration of 24.5 years (Q1; Q3 17.0; 42.5) and a median SELENA-SLEDAI score of 7.0 (Q1; Q3 5.3; 15.3). The median age of healthy controls was 33.5 years (Q1; Q3 25.0; 43.5). The concentration of MP DNA positively correlated with anti-dsDNA IgG ($r[10] = 0.806$, $P = 0.005$) (Fig. 3) after exclusion of outliers, which may be consistent with MP DNA as an antigenic source for anti-dsDNA IgG formation. However, MP DNA did not correlate with complement levels or SELENA-SLEDAI scores.

The results from this study provide new perspectives on the presence of extracellular DNA in SLE patients and healthy controls. Using flow cytometry, MP DNA were visualised and enumerated in the PFP of SLE patients and healthy controls. In the form of MPs, DNA likely exists on both the surface and interior.¹⁰ This would account for the observation of DNA packaged within MPs being protected from DNase activity.¹¹ Further, plasma DNA not associated with MPs were detected in both SLE patients and healthy controls. Experiments to assess the origins of plasma DNA and its sensitivity to DNase I were not

performed in the current study. The most striking finding in this study is the marked difference in the concentration of MP DNA in SLE patients compared to healthy controls. Although the increased MP numbers suggest a role in SLE immunopathogenesis, elucidating this role is difficult at present. Ullal et al showed some increase in the total number of MP and increase in IgG positive MPs in SLE patients compared to healthy controls.¹² Nielsen et al showed that MP populations that do not bind annexin V are increased whereas total number of MP and annexin V-binding MPs are decreased in SLE patients compared to healthy controls.¹³ We showed a correlation between MP DNA concentration and anti-dsDNA IgG titers. Ullal et al and Nielsen et al showed a correlation between the number of IgG-positive MPs and anti-DNA levels in SLE patients.^{12,14} However, their findings did not demonstrate that the bound IgG antibody was anti-dsDNA IgG. In this regard, IgG positive MPs can also occur in the synovial fluid of rheumatoid arthritis patients.^{12,15} Hence, to our knowledge, this is the first study to demonstrate the presence of MP DNA definitively in the




Fig. 2. Concentration of MP DNA in SLE patients and healthy controls. The median MP DNA/ μ L of plasma for 14 SLE patients was significantly higher than the 8 healthy controls.




Fig. 3. The relationship between MP DNA and anti-dsDNA levels in the plasma of SLE patients. There was significant correlation of MP DNA concentration with anti-dsDNA IgG titers.

plasma of SLE patients. The correlation with anti-dsDNA IgG suggests that MP DNA may be a source of immunogenic autoantigen for the production of anti-dsDNA antibodies and our results support this idea. Further studies will refine the role of MP DNA in the pathogenesis, perpetuation and modulation of disease activity in SLE.

REFERENCES

- Celada F. The anti-DNA story. *Lupus* 2011;20:553-5.
- Pisetsky DS. The origin and properties of extracellular DNA: from PAMP to DAMP. *Clin Immunol* 2012;144:32-40.
- Rekvi OP, Mortensen ES. Immunity and autoimmunity to dsDNA and chromatin—the role of immunogenic DNA-binding proteins and nuclease deficiencies. *Autoimmunity* 2012;45:588-92.
- Rekvi OP, Nossent JC. Anti-double-stranded DNA antibodies, nucleosomes, and systemic lupus erythematosus: a time for new paradigms? *Arthritis Rheum* 2003;48:300-12.
- Mause SF, Weber C. Microparticles: protagonists of a novel communication network for intercellular information exchange. *Circ Res* 2010;107:1047-57.
- Halicka HD, Bedner E, Darzynkiewicz Z. Segregation of RNA and separate packaging of DNA and RNA in apoptotic bodies during apoptosis. *Exp Cell Res* 2000;260:248-56.
- Pisetsky DS, Ullal AJ, Gauley J, Ning TC. Microparticles as mediators and biomarkers of rheumatic disease. *Rheumatology (Oxford)* 2012;51:1737-46.
- Hochberg MC. Updating the American College of Rheumatology revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1997;40:1725.
- Petri M, Kim MY, Kalunian KC, Grossman J, Hahn BH, Sammaritano LR, et al. Combined oral contraceptives in women with systemic lupus erythematosus. *N Engl J Med* 2005;353:2550-8.
- Pisetsky DS, Lipsky PE. Microparticles as adjuvants in the pathogenesis of SLE. *Nat Rev Rheumatol* 2010;6:368-72.
- Orozco AF, Jorgez CJ, Horne C, Marquez-Do DA, Chapman MR, Rodgers JR, et al. Membrane protected apoptotic trophoblast microparticles contain nucleic acids: relevance to preeclampsia. *Am J Pathol* 2008;173:1595-608.
- Ullal AJ, Reich CF 3rd, Clowse M, Criscione-Schreiber LG, Tochacek M, Monestier M, et al. Microparticles as antigenic targets of antibodies to DNA and nucleosomes in systemic lupus erythematosus. *J Autoimmun* 2011;36:173-80.
- Nielsen CT, Ostergaard O, Johnsen C, Jacobsen S, Heegaard NH. Distinct features of circulating microparticles and their relationship to clinical manifestations in systemic lupus erythematosus. *Arthritis Rheum* 2011;63:3067-77.
- Nielsen CT, Ostergaard O, Stener L, Iversen LV, Truedsson L, Gullstrand B, et al. Increased IgG on cell-derived plasma microparticles in systemic lupus erythematosus is associated with autoantibodies and complement activation. *Arthritis Rheum* 2012;64:1227-36.
- Cloutier N, Tan S, Boudreau LH, Cramb C, Subbiah R, Lahey L, et al. The exposure of autoantigens by microparticles underlies the formation of potent inflammatory components: the microparticle-associated immune complexes. *EMBO Mol Med* 2013;5:235-49.

Sen Hee Tay,^{1,2} MBBS, M Med, MRCP, Aisha Lateef,^{1,2} MBBS, MRCP, FAMS, Yaw Chyn Lim,³ PhD, Irene JV Ng,² BSc, Dow Rhon Koh,^{2,4} MBBS, M Med, PhD, Lina HK Lim,⁴ PhD

¹Department of Medicine, Yong Loo Lin School of Medicine, National University of Singapore, Singapore

²Division of Rheumatology, Department of Medicine, National University Hospital, National University Health System, Singapore

³Department of Pathology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore

⁴Department of Physiology & NUS Immunology Programme, Yong Loo Lin School of Medicine, National University of Singapore, Singapore

Address for Correspondence: Dr Tay Sen Hee, Division of Rheumatology, Department of Medicine, National University Hospital, National University Health System, Level 10, NUHS Tower Block, 1E Kent Ridge Road, Singapore 119228.

Email: sen_hee_tay@nuhs.edu.sg