

Supplementary Table 1. Allelic and genotypic frequencies and Hardy-Weinberg equilibrium of the study population.

Genetic polymorphisms	Allelic frequencies		Genotypic frequencies (n, %)			<i>P</i> value for HWE
	Wild-type allele	Variant allele	Wild-type homozygotes	Heterozygotes	Variant homozygotes	
<i>F2</i> rs5896 (C>T)	0.40	0.60	68 (14.7)	237 (51.1)	159 (34.3)	0.18
<i>F10</i> rs5960 (C>T)	0.48	0.52	103 (22.2)	243 (52.4)	118 (25.4)	0.30
<i>FGA</i> rs2070011 (T>C)	0.56	0.44	143 (30.6)	233 (49.9)	91 (19.5)	0.82
<i>FGA</i> rs6050 (T>C)	0.47	0.53	103 (22.1)	231 (49.5)	133 (28.5)	0.89
<i>FGA</i> rs2070022 (G>A)	0.87	0.13	352 (75.7)	107 (23)	6 (1.3)	0.50
<i>FGB</i> rs1800788 (C>T)	0.35	0.65	58 (12.5)	208 (44.8)	198 (42.7)	0.77
<i>FGB</i> rs4220 (G>A)	0.86	0.14	341 (73)	117 (25.1)	9 (1.9)	0.78
<i>FGB</i> rs4463047 (T>C)	0.49	0.51	105 (23)	235 (51.5)	116 (25.4)	0.50
<i>FGG</i> rs2066865 (G>A)	0.47	0.53	103 (22.1)	233 (50)	130 (27.9)	0.94
<i>FGG</i> rs1800792 (T>C)	0.69	0.31	220 (47.2)	201 (43.1)	45 (9.7)	0.93

HWE: Hardy-Weinberg equilibrium.