

Table 1. Evaluated parameters at consecutive follow-up visits.

	<b>Pre-MPTCP</b>	<b>Day 1</b>	<b>Week 1</b>	<b>Week 2</b>	<b>Month 1</b>	<b>Month 3</b>	<b>Month 6</b>	<b>Month 12</b>
No. of eyes that underwent further intervention	-	0	0	0	1	4	5	5
No. of eyes with complications	-	0	0	0	0	0	0	0
IOP, mmHg <sup>a</sup>	27.4 (2.3)	19.7 (2.9)	15.7 (4.2)	24.0 (2.7)	24.7 (2.7)	24.3 (3.0)	20.1 (3.0)	23.0 (3.3)
Treatment success, %	-	55.6	75.0	60.0	50.0	50.0	66.7	28.6
No. of glaucoma medications <sup>a</sup>	1.3 (0.1)	1.2 (0.1)	1.2(0.1)	1.2 (0.1)	1.2 (0.1)	1.3 (0.1)	1.2 (0.1)	1.2 (0.1)
No. of eyes requiring acetazolamide	2	3	2	2	1	0	0	0
VA, LogMAR <sup>a</sup>	0.81 (0.24)	0.96 (0.24)	1.05 (0.25)	1.01 (0.24)	1.02 (0.24)	1.05 (0.24)	1.08 (0.24)	1.12 (0.25)
MD of HVF, dB <sup>a</sup>	-19.30 (3.03)	-	-	-	-	-	-	-18.90 (2.63)
Duration of postoperative steroids eye drops, days <sup>b,c</sup>	-	-	-	-	-	-	-	17.5 (7–365)
Duration of follow-up, months <sup>c</sup>	-	-	-	-	-	-	-	7.5 (0.6–11.4)

dB: decibels; HVF: Humphrey visual field; IOP: intraocular pressure; LogMAR: logarithm of minimum angle of resolution; MPTCP: micropulse transscleral cyclophototherapy; MD: mean deviation; VA: visual acuity

<sup>a</sup> Marginal means (standard error).

<sup>b</sup>  $P < 0.05$  when compared with parameters before micropulse transscleral cyclophototherapy.

<sup>c</sup> Median (minimum–maximum).