

Supplementary Table S3: Summary of reported Hantavirus cases in Singapore.

	Year of diagnosis	Age/Sex	Past medical history	Clinical presentation, occupation and travel history	Complications	Diagnostic method	Clinical outcomes
Chan, et al. 1996	1996	67/Male	Diabetes mellitus, hypertension	Fever, lethargy and confusion. Ragman. No documented travel history.	Acute kidney injury, thrombocytopenia, severe liver dysfunction, hypotension, stroke	Specific antibody testing with immunofluorescence assay	Survived, but prolonged hospital stay (5 months). Good renal recovery. Ambulating with walking frame on discharge.
Chan, et al. 2013	2010	42/Male	None	Fever, myalgia, lethargy for 3 days. Construction site supervisor. From Johor Bahru, Malaysia.	Acute kidney injury, thrombocytopenia, severe liver dysfunction.	Hantavirus serology (EIA)	Survived, with good functional and renal recovery.
Chan, et al. 2013	2010	35/Male	None	Fever, myalgia, nausea/vomiting and abdominal pain for 1 week. Office worker. Frequent travel to Johor Bahru, and Kuala Lumpur, Malaysia.	Acute kidney injury, thrombocytopenia, mild transaminitis	Hantavirus serology (EIA)	Survived, with good functional and renal recovery
Ngiam, et al. (current case)	2022	59/Male	None	Fever, lethargy, myalgia, unsteady gait for 1 week. Shipyard engineer. Frequent travel to Ipoh, Malaysia.	Acute kidney injury, thrombocytopenia, mild transaminitis.	Hantavirus serology (EIA), followed by specific antibody testing with immunofluorescence assay	Survived, with good functional and renal recovery