

Variables	Alteplase dose (mg) ^a				P value ^b	Total ^c n=129
	<2.5 n=17	2.5 n=36	5 n=61	10 n=12		
Age					0.29	
Mean (SD)	69.5 (13.4)	67.1 (17.3)	62.8 (14.9)	62.6 (16.1)		65.1 (15.5)
Median (minimum, maximum)	69 (47, 95)	66.5 (19, 97)	62 (22, 89)	66 (30, 80)		66 (19, 97)
Sex, no. (%)					0.35	
Female	3 (17.7)	12 (33.3)	13 (21.3)	1 (8.3)		30 (23.3)
Male	14 (82.3)	24 (66.7)	48 (78.7)	11 (91.7)		99 (76.7)
BMI					0.18	
No.	17	32	51	9		112
Mean (SD)	23.2 (4.8)	20.3 (4.7)	22.2 (4.7)	20.6 (3.6)		21.6 (4.7)
Median (minimum, maximum)	22.6 (17.6, 36.2)	20.3 (13.2, 35.7)	21.4 (13.8, 36.2)	20.7 (16.1, 27.9)		21.1 (13.2, 36.2)
RAPID score					0.1	
Mean (SD)	4.2 (1.3)	3.5 (1.1)	3.4 (1.1)	3.5 (0.7)		3.5 (1.1)
Median (minimum, maximum)	4 (2, 6)	3 (2, 6)	3 (2, 6)	3 (3, 5)		3 (2, 6)

BMI: body mass index; RAPID score: renal, age, purulence, infection source and dietary factor; SD: standard deviation

^a Three subjects had no record of alteplase dose.

^b P value was obtained using one-way analysis of variance or Fisher's Exact test (sex).

^c All 129 subjects were included.