

Variables	Alteplase dose (mg)				Total
	<2.5 n=17	2.5 n=37	5 n=62	10 n=12	
Baseline CXR loculation, no. (%)					
Yes	4 (23.5)	14 (37.8)	16 (25.8)	4 (33.3)	39 (29.8)
No	13 (76.5)	23 (62.2)	46 (74.2)	8 (66.7)	92 (70.2)
Loculations on CT, no. (%)					
Yes	9 (52.9)	23 (65.7)	48 (77.4)	7 (58.3)	89 (69)
No	8 (47.1)	12 (34.3)	14 (22.6)	5 (41.7)	40 (31)
Pleural thickening >2 mm, no. (%)					
Yes	13 (76.5)	24 (68.6)	46 (75.4)	12 (100)	96 (75)
No	4 (23.5)	11 (31.4)	15 (24.6)	0	32 (25)
Time from chest tube insertion to first dose (days)					
Mean (SD)	13 (8.3)	12.6 (13.8)	8.5 (11.5)	8.3 (7.4)	10.2 (11.5)
Median (minimum, maximum)	10.5 (2, 27)	7 (1, 60)	5 (1, 67)	6 (1, 21)	6.5 (1, 67)

CXR: chest X-ray; CT: computed tomography; SD: standard deviation