

Demographics	Orthogeriatric care cohort (n=252)
Age, mean (SD)	77.7 (10.0)
Female, no. (%)	204 (81.0)
CFS group, no. (%)	
1–3	55 (21.8)
4–9	197 (78.2)
Location of fracture, no. (%)	
Femur	77 (30.6)
Patella	30 (11.9)
Tibia	33 (13.1)
Ankle	36 (14.3)
Acetabulum	20 (7.9)
Periprosthetic, peri-implant	56 (22.2)
Mobility on admission, no. (%)	
Community ambulant	147 (58.3)
Homebound	105 (41.7)
Level of assisted mobility, no. (%)	
Independent of aid	133 (52.8)
Single arm aid	45 (17.9)
Double arm aid	15 (5.9)
Personal assistance	32 (12.7)
Wheelchair/bedbound	27 (10.7)
Treatment type, no. (%)	
Conservative	108 (42.9)
Operated	144 (57.1)
Weight-bearing status post-intervention, no. (%)	
Non-weight bearing	143 (56.7)
Partial weight bearing (i.e. 50% of weight bearing on affected limb)	23 (9.1)
Weight bearing as tolerated	89 (35.3)
Outcomes	
Length of stay, days; mean (SD)	
Conservative	8.6 (9.9)

Demographics	Orthogeriatric care cohort (n=252)
Operative	10.9 (7.7)
Total	9.4 (6.5)
Time to surgery, days; mean (SD)	2.9 (2.5)
Inpatient medical complications, no. (%)	
Urinary tract infection	18 (7.1)
Pneumonia	6 (2.4)
Delirium	13 (5.2)
DVT/PE	7 (2.8)
Stroke	3 (1.2)
Acute myocardial infarction	0 (0)
Gastrointestinal bleeding	3 (1.2)
Total incidence	50 (17.9)
Inpatient mortality, no. (%)	5 (2.0)
One-year mortality, no. (%)	28 (11.1)

CFS: Clinical Frailty Score; DVT: deep vein thrombosis;
PE: pulmonary embolism; SD: standard deviation